

Technical data

Sildenafil 50

Sildenafil 50mg coated tablets



Drug Name	Sildenafil 50
Generic name	Sildenafil 50 mg coated tablets
Qualitative-Quantitative Composition	Each coated tablet contains: Sildenafil Citrate 70.24 mg equivalent to Sildenafil 50mg Excipients q.s.
Pharmaceutical form	Coated Tablets
Clinical Data	
- Therapeutic indications	Treatment of erectile dysfunction in men, which is the inability to obtain or maintain an erection sufficient for satisfactory sexual activity. For sildenafil to be effective, sexual stimulation is necessary.
- Route of Administration	Oral
- Contraindications	Hypersensitivity to the active substance or to any of the excipients. Consistent with its known effects on the nitric oxide/cyclic guanosine monophosphate (cGMP) pathway, sildenafil has been shown to potentiate the hypotensive effects of nitrates, and concomitant administration of sildenafil with nitric oxide donors (such as amyl nitrite) is contraindicated. or nitrates in any of their forms. Drugs for the treatment of erectile dysfunction, including sildenafil, should not be used in men for whom sexual activity is inadvisable (for example: patients with serious cardiovascular dysfunctions such as unstable angina or severe heart failure). Sildenafil is contraindicated in patients who have lost vision in one eye due to non-arteritic anterior ischemic optic neuropathy (NAION), regardless of whether or not this event was associated with prior exposure to a PDE5 inhibitor.
- Presentation	Blister of 4 coated tablets in a box.
- Incompatibilities	Does not apply
prescription regime	<i>Use in adults:</i> The recommended dose is 50 mg taken as needed, approximately one hour before sexual activity. Based on efficacy and tolerance, the dose may be increased to 100 mg or decreased to 25 mg. The maximum recommended dose is 100 mg. The maximum recommended dosage schedule is once daily. If Sildenafil is taken with food, the onset of its pharmacological action may be delayed compared to when taken on an empty stomach. <i>Use in the elderly:</i> No dose adjustment is required in the elderly. <i>Use in patients with renal insufficiency:</i> The recommended doses described in "Use in adults" are applicable to patients with mild to moderate renal insufficiency (creatinine clearance = 30-80 ml/min).
Storage Condition	Store at a temperature below 30°C. Protect from light and moisture. Keep out of the reach of children.
special legends	Does not apply
Countries where it is approved	Honduras, El Salvador, Costa Rica, Guatemala
Laboratory Manufacturer	Saga Lifesciences Limited, Ahmedabad, India
Review date	September 2020